

Zakrzewicz A, Richter K, Agné A, Wilker S, Siebers K, Fink B, Krasteva-Christ G, Althaus M, Padberg W, Hone AJ, McIntosh JM, Grau V.

[Canonical and Novel Non-Canonical Cholinergic Agonists Inhibit ATP-Induced Release of Monocytic Interleukin-1 \$\beta\$ via Different Combinations of Nicotinic Acetylcholine Receptor Subunits \$\alpha 7\$, \$\alpha 9\$ and \$\alpha 10\$.](#)

Frontiers in Cellular Neuroscience 2017, 11, 189.

Copyright:

© 2017 Zakrzewicz, Richter, Agné, Wilker, Siebers, Fink, Krasteva-Christ, Althaus, Padberg, Hone, McIntosh and Grau. This is an open-access article distributed under the terms of the [Creative Commons Attribution License \(CC BY\)](#). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

DOI link to article:

<https://doi.org/10.3389/fncel.2017.00189>

Date deposited:

28/07/2017



This work is licensed under a [Creative Commons Attribution 4.0 International License](#)



Canonical and Novel Non-Canonical Cholinergic Agonists Inhibit ATP-Induced Release of Monocytic Interleukin-1 β via Different Combinations of Nicotinic Acetylcholine Receptor Subunits $\alpha 7$, $\alpha 9$ and $\alpha 10$

OPEN ACCESS

Edited by:

Barbara Jane Morley,
Boys Town National Research
Hospital, United States

Reviewed by:

Richard J. Lewis,
The University of Queensland,
Australia
Koichiro Kawashima,
Kitasato University, Japan

*Correspondence:

Anna Zakrzewicz
anna.zakrzewicz@chiru.med.uni-
giessen.de
Katrin Richter
katrin.richter@chiru.med.uni-
giessen.de

[†]These authors have contributed
equally to this work.

Received: 25 April 2017

Accepted: 20 June 2017

Published: 05 July 2017

Citation:

Zakrzewicz A, Richter K, Agné A,
Wilker S, Siebers F, Fink B,
Krasteva-Christ G, Althaus M,
Padberg W, Hone AJ, McIntosh JM
and Grau V (2017) Canonical and
Novel Non-Canonical Cholinergic
Agonists Inhibit ATP-Induced Release
of Monocytic Interleukin-1 β via
Different Combinations of Nicotinic
Acetylcholine Receptor Subunits $\alpha 7$,
 $\alpha 9$ and $\alpha 10$.
Front. Cell. Neurosci. 11:189.
doi: 10.3389/fncel.2017.00189

Anna Zakrzewicz^{1*†}, Katrin Richter^{1*†}, Alisa Agné¹, Sigrid Wilker¹, Kathrin Siebers¹,
Bijan Fink¹, Gabriela Krasteva-Christ^{2,3,4}, Mike Althaus^{5,6}, Winfried Padberg¹,
Arik J. Hone⁷, J. Michael McIntosh^{7,8,9} and Veronika Grau^{1,4}

¹Laboratory of Experimental Surgery, Department of General and Thoracic Surgery, Justus-Liebig-University Giessen, Giessen, Germany, ²Institute of Anatomy and Cell Biology, Justus-Liebig-University Giessen, Giessen, Germany, ³Institute of Anatomy and Cell Biology, Saarland University, Homburg, Germany, ⁴Member of the German Centre for Lung Research, Giessen, Germany, ⁵Institute of Animal Physiology, Justus-Liebig-University Giessen, Giessen, Germany, ⁶School of Biology, Newcastle University, Newcastle upon Tyne, United Kingdom, ⁷Department of Biology, University of Utah, Salt Lake City, UT, United States, ⁸George E. Wahlen Veterans Affairs Medical Center, Salt Lake City, UT, United States, ⁹Department of Psychiatry, University of Utah, Salt Lake City, UT, United States

Recently, we discovered a cholinergic mechanism that inhibits the adenosine triphosphate (ATP)-dependent release of interleukin-1 β (IL-1 β) by human monocytes via nicotinic acetylcholine receptors (nAChRs) composed of $\alpha 7$, $\alpha 9$ and/or $\alpha 10$ subunits. Furthermore, we identified phosphocholine (PC) and dipalmitoylphosphatidylcholine (DPPC) as novel nicotinic agonists that elicit metabotropic activity at monocytic nAChR. Interestingly, PC does not provoke ion channel responses at conventional nAChRs composed of subunits $\alpha 9$ and $\alpha 10$. The purpose of this study is to determine the composition of nAChRs necessary for nicotinic signaling in monocytic cells and to test the hypothesis that common metabolites of phosphatidylcholines, lysophosphatidylcholine (LPC) and glycerophosphocholine (G-PC), function as nAChR agonists. In peripheral blood mononuclear cells from nAChR gene-deficient mice, we demonstrated that inhibition of ATP-dependent release of IL-1 β by acetylcholine (ACh), nicotine and PC depends on subunits $\alpha 7$, $\alpha 9$ and $\alpha 10$. Using a panel of nAChR antagonists and siRNA technology, we confirmed the involvement of these subunits in the control of IL-1 β release in the human monocytic cell line U937. Furthermore, we showed that LPC (C16:0) and G-PC efficiently inhibit ATP-dependent release of IL-1 β . Of note, the inhibitory effects mediated by LPC and G-PC depend on nAChR subunits $\alpha 9$ and $\alpha 10$, but only to a small degree on $\alpha 7$. In *Xenopus laevis* oocytes heterologously expressing different combinations of human $\alpha 7$, $\alpha 9$ or $\alpha 10$ subunits, ACh induced canonical ion channel activity, whereas LPC, G-PC and PC did not. In conclusion,

we demonstrate that canonical nicotinic agonists and PC elicit metabotropic nAChR activity in monocytes via interaction of nAChR subunits $\alpha 7$, $\alpha 9$ and $\alpha 10$. For the metabotropic signaling of LPC and G-PC, nAChR subunits $\alpha 9$ and $\alpha 10$ are needed, whereas $\alpha 7$ is virtually dispensable. Furthermore, molecules bearing a PC group in general seem to regulate immune functions without perturbing canonical ion channel functions of nAChR.

Keywords: acetylcholine, *CHRNA*, glycerophosphocholine, inflammasome, interleukin-1beta, lysophosphatidylcholine, nicotine and phosphocholine

INTRODUCTION

Interleukin-1 β (IL-1 β) is a potent pro-inflammatory cytokine of innate immunity that importantly contributes to host defense against pathogens (Vladimer et al., 2013). Production and secretion of IL-1 β are strictly controlled and typically require two independent danger signals (Rathinam et al., 2012). The presence of potential pathogens is commonly sensed by Toll-like receptors that lead to cytoplasmic expression of pro-IL-1 β . A prototypical second signal is extracellular adenosine triphosphate (ATP) originating from the cytoplasm of damaged host cells (Trautmann, 2009). ATP binds to the ionotropic P2X7 receptor, which enables efflux of potassium ions. This induces the assembly of the NLRP3 (NLR family, pyrin domain containing protein 3) inflammasome and activation of the protease caspase-1, which eventually cleaves pro-IL-1 β and enables the release of mature bioactive IL-1 β (Ferrari et al., 2006; Gross et al., 2011; Ozaki et al., 2015). Excessive systemic release of IL-1 β can be dangerous for the host (Dinarello et al., 2012). Elevated systemic levels of IL-1 β can cause fever, shock, systemic inflammatory response syndrome (SIRS) and can lead to life-threatening multi-organ damage (Cauwels et al., 2014). Mechanisms controlling IL-1 β maturation are of substantial clinical interest but so far have remained largely unexplored.

Recently, we demonstrated that activation of nicotinic acetylcholine (ACh) receptors (nAChRs) by classical ligands like ACh, nicotine (Nic) and choline (Cho) inhibits ATP signaling (Hecker et al., 2009, 2015; Mikulski et al., 2010; Richter et al., 2016), activation of caspase-1 and release of IL-1 β by human monocytes (Hecker et al., 2015; Richter et al., 2016). Interestingly, activated monocytes, T cells and vascular endothelial cells can endogenously produce ACh, which presumably acts in an autocrine or paracrine fashion (Hecker et al., 2009; Wilczynska et al., 2011; Kawashima et al., 2012). In addition to conventional nAChR agonists, we demonstrated that free phosphocholine (PC), a common metabolite of phosphatidylcholine and PC-modified macromolecules can also effectively inhibit the ATP-dependent release of IL-1 β by human and murine monocytes (Hecker et al., 2015; Richter et al., 2016). Gene-silencing experiments and pharmacological studies demonstrated that nAChR subunits $\alpha 7$, $\alpha 9$ and/or $\alpha 10$ are involved in the cholinergic inhibition of IL-1 β release (Hecker et al., 2015). The nAChR subunits $\alpha 9$ and $\alpha 10$ contribute

to the inhibitory effect of PC (Richter et al., 2016). It is however unclear, if the effect of PC also depends on subunit $\alpha 7$ and if receptor subunits $\alpha 7$, $\alpha 9$ and $\alpha 10$ cooperate in monocytic cells or exert redundant functions. Similarly, dipalmitoylphosphatidylcholine (DPPC), the main lipid component of pulmonary surfactant, inhibits ATP-driven inflammasome activation in monocytes. DPPC signals via the nAChR subunit $\alpha 9$ in combination with either subunit $\alpha 7$ or $\alpha 10$ (Backhaus et al., 2017).

Phosphatidylcholines possess a PC head group and two non-polar fatty acid chains. Enzymatic removal of one fatty acid results in lysophosphatidylcholine (LPC), which can be further metabolized to glycerophosphocholine (G-PC). LPC has been shown to modulate responses of the innate and adaptive immune system by various non-cholinergic mechanisms (Kabarowski et al., 2002; Stock et al., 2006; Carneiro et al., 2013). The potential involvement of LPC and G-PC, two molecules bearing a PC head group, in the cholinergic regulation of ATP-mediated IL-1 β release by monocytes has not yet been tested.

The purpose of this study was to investigate if nAChR subunits $\alpha 7$, $\alpha 9$ and $\alpha 10$ interact in monocytes and which nAChR subunit composition is necessary for ACh-, Nic- and PC-mediated inhibition of IL-1 β release. Furthermore, we hypothesize that LPC and G-PC, both PC-bearing metabolites of phosphatidylcholines, can function as agonists of nAChRs in monocytes and contribute to the regulation of IL-1 β release. In addition, we characterize, which nAChR subunits are required for the inhibitory effect mediated by LPC and G-PC. Finally, we investigate if LPC, G-PC and PC induce ionotropic functions at heterologously expressed canonical nAChRs.

MATERIALS AND METHODS

Animals

This study was carried out in accordance with the recommendations of the NIH "Guide for the Care and Use of Laboratory Animals". The protocols for animal experiments were approved by the Regierungspräsidium Giessen, Hesse, Germany (permit number 549_M; R.P. Nr. Gi 20/23-Nr. A12/2011; R.P. Nr. Gi 20/23-Nr. A10/2011). Adult female *Xenopus laevis* (Xenopus-Express, Le Bourg, France) as well as male and female wild-type (WT) and *Chrna7* (C57B1/6J), *Chrna9* (129S-*Chrna9*^{tm1Bdv}/J) and *Chrna10* (129S4-*Chrna10*^{tm1Bdv}/Mmucd) gene-deficient mice

were used. The generation and characterization of every gene-deficient mouse strain was described before (Orr-Urtreger et al., 1997; Vetter et al., 1999, 2007; Whiteaker et al., 2007). *Chrna 7* gene-deficient mice were obtained from the Jackson Laboratories after being re-derived, whereas *Chrna 9* and *Chrna 10* gene-deficient mice were kindly provided by Prof. Douglas E. Vetter. The genotype of every mouse was evaluated by PCR. The corresponding WT background strain for every gene-deficient mouse was used as a control.

Mononuclear Leukocytes from Mice

Mouse peripheral blood mononuclear cells (PBMCs) were freshly isolated from heparinized blood obtained from WT, *Chrna7*, *Chrna9*, or *Chrna10* gene-deficient mice using Percoll (GE Healthcare Bio-Sciences AB, Uppsala, Sweden; 1.082 g/ml) density gradient centrifugation. Mononuclear cells were cultured for 2 h in 24-well-plates at 37°C, 5% CO₂, in RPMI 1640 (Gibco/Life Technologies, Carlsbad, CA, USA) supplemented with 10% fetal bovine serum (FBS Superior EU, Biochrom GmbH, Berlin, Germany) and 2 mM L-glutamine (GlutaMAXTM, Gibco/Life Technologies). Thereafter, cells were stimulated with 2'(3')-O-(4-benzoylbenzoyl)adenosine 5'-triphosphate triethylammonium salt (BzATP, 100 μ M, Sigma-Aldrich, Taufkirchen, Germany) for 30 min, in the presence or absence of ACh chloride (10 μ M), PC chloride calcium salt tetrahydrate (PC, 100 μ M) or nicotine (Nic, 100 μ M; all purchased from Sigma-Aldrich, Taufkirchen, Germany). Cell culture supernatants were collected and stored at -20°C until IL-1 β and lactate dehydrogenase (LDH) measurement.

U937 Cells

The human histocytic lymphoma cell line U937 was obtained from the German Collection of Microorganisms and Cell Cultures (Braunschweig, Germany) and cultured according to the protocol described for mouse PBMCs. Cells in the log-phase of growth were transferred to 24-well plates (1 \times 10⁶ cells/ml and per well) and primed with 1 μ g/ml LPS from *Escherichia coli* (L2654; Sigma-Aldrich) for 5 h. Thereafter, BzATP (100 μ M) was applied for 30 min, in the presence or absence of ACh (10 μ M), Nic (100 μ M), PC (100 μ M), L- α -G-PC (Sigma-Aldrich; 0.1 μ M to 1000 μ M) or 1-palmitoyl-*sn*-glycero-3-phosphocholine LPC (C16:0), Sigma-Aldrich; 0.1 μ M to 100 μ M). To test the involvement of different subunits of nAChR, the following antagonists were applied: mecamylamine hydrochloride (Mec, 100 μ M, Sigma-Aldrich), α -bungarotoxin (α -Bun, 1 μ M, Tocris Bioscience, Bristol, UK), strychnine hydrochloride (Stry, 10 μ M, Sigma-Aldrich) or conotoxins ArIB [V11L, V16D] (500 nM) or RgIA4 (200 nM; Ramarao and Cohen, 1998; Whiteaker et al., 2007; Innocent et al., 2008; Richter et al., 2016; Romero et al., 2017).

Knock-Down of $\alpha 7$, $\alpha 9$ or $\alpha 10$ nAChR Subunit Expression

The expression of human nAChR subunits $\alpha 7$, $\alpha 9$ or $\alpha 10$ was attenuated using siRNA technology. Cells were transfected with ON-TARGETplus human *CHRNA7* ($\alpha 7$), *CHRNA9* ($\alpha 9$) or *CHRNA10* ($\alpha 10$) SMARTpool siRNA (Thermo Fisher Scientific,

Schwerte, Germany) or with ON-TARGETplus non-targeting control pool (Thermo Fisher Scientific) as a negative control. Cells were transfected with 30 pmol siRNA/1 \times 10⁶ cells using the Amaxa Cell Line Nucleofector Kit C and the Nucleofector II Device (both from Lonza, Cologne, Germany) according to the manufacturer's instructions. The siRNA-mediated down-regulation of target genes was assessed 48 h after transfection by real-time RT-PCR.

RNA Isolation and cDNA Synthesis

Total RNA was isolated from 1 \times 10⁶ U937 cells using the RNeasy Plus Mini Kit (Qiagen, Hilden, Germany) according to the manufacturer's instructions. One microgram of RNA was reversely transcribed using M-MLV H⁻ Reverse Transcriptase and 1 μ g of random hexamer primers (Promega, Mannheim, Germany).

Real-Time RT-PCR

To confirm efficient and selective knock-down of nAChR subunits $\alpha 7$, $\alpha 9$ or $\alpha 10$ targeted selectively with siRNA, real-time RT-PCR was performed ($n = 4$ per experimental group, each sample was assessed in duplicates) using the ABI 7900 Sequence Detection System (Applied Biosystems, Foster City, CA, USA) and Platinum SYBER green qPCR Super Mix-UDG (Invitrogen, Karlsruhe, Germany). The hydroxymethylbilane synthase (HMBS) gene was selected as a reference gene, as it was reported not to be regulated in monocytes under various culture conditions and stimulations (Moosig et al., 2006). Primers specific for the detection of HMBS, $\alpha 7$, $\alpha 9$ and $\alpha 10$ nAChR subunits were synthesized by MWG Biotech (Ebersberg, Germany) and their sequences have been published before (Hecker et al., 2015). Changes in the mRNA expression levels of nAChR were calculated by the 2 $^{\Delta\Delta CT}$ method, where ΔCT represents the difference between the CT value of HMBS gene and the CT value of gene of interest. The mean of the mRNA expression values from cells transfected with non-targeting (scrambled) siRNA was set to one and the values from cells transfected with siRNA specific for nAChR were calculated accordingly.

ELISA

The IL-1 β concentration in the cell culture supernatants was determined by mouse- or human- specific Quantikine Immunoassays (R&D Systems, Minneapolis, MN, USA) according to the manufacturer's instructions.

LDH Measurement

The viability of the cells was estimated by measurement of LDH in cell culture supernatants by the Non-Radioactive Cytotoxicity Assay (Promega, Madison, WI, USA) according to the manufacturer's instructions. To assess the proportion of dead cells, a maximum LDH release control was generated. For this purpose, an equivalent number of U937 cells were disrupted by freezing them twice at -80°C. The LDH value of the sample of interest was compared with the total content of LDH in the control sample, which was set to 100% and the relative cell death was calculated accordingly.

Isolation and Culture of Oocytes

Xenopus laevis oocytes were purchased from Ecocyte Bioscience (Castrop-Rauxel, Germany) or obtained from adult female frogs (Xenopus-Express, Le Bourg, France) and prepared as previously described (Richter et al., 2016). Defolliculated oocytes were stored in oocyte Ringer's solution (ORi) containing (in mM) 90 NaCl, 1 KCl, 2 CaCl₂, 5 HEPES (4-(2-hydroxyethyl)-piperazine-1-ethanesulfonic acid), 2.5 pyruvate, 20 mg/ml penicillin and 25 mg/ml streptomycin (pH 7.4) at 16°C. All chemicals used for ORi preparation were purchased from Fluka (Deisenhofen, Germany), except for HEPES, penicillin and streptomycin (Sigma-Aldrich).

Heterologous Expression of Human nAChRs in Oocytes and Electrophysiological Whole-Cell Measurements

The cDNA clones of human α 9, human α 10 and human 43 kDa receptor-associated protein of the synapse (RAPSIN) in the pTNT vector were obtained from Eurofins Genomics (Ebersberg, Germany). Capped cRNA was synthesized using an *in vitro* transcription kit (T7 RiboMAXTM Large Scale RNA Production System Kit, Promega, Mannheim, Germany). The cRNA encoding human α 7 was kindly provided by G. Schmalzing (Department of Molecular Pharmacology, RWTH Aachen University, Aachen, Germany). In brief, a plasmid containing the entire coding region of human α 7 (GenBank NM_000746.5, Addgene plasmid #62276; Wang et al., 2014) was purchased from Addgene (Teddington, UK¹) and subcloned via gateway PCR from the pcDNA3.1 vector into a gateway-modified version of the oocyte expression vector pNKS2 (Gloor et al., 1995). The plasmid was linearized downstream of the vector-provided polyA tail with XhoI and transcribed with SP6 RNA polymerase into capped and polyadenylated cRNA, as previously described (Stolz et al., 2015).

Oocytes were injected with cRNA encoding nAChR subunit α 7 (20 ng/oocyte) or with cRNA combinations for subunits α 7 and α 9 (20 ng/oocyte, each), or subunits α 9 and α 10 (20 ng/oocyte, each), or α 7, α 9 and α 10 (16, 19, 19 ng/oocyte, respectively). RAPSIN (5 ng/oocyte) was co-injected in all experiments to increase the expression levels and to obtain stable nAChR expression (Froehner et al., 1990). The cRNA was dissolved in nuclease-free water and injected in a volume of 50.6 nl per oocyte. In all two-electrode voltage-clamp (TEVC) experiments representative controls were performed with oocytes that were injected with the same amount of nuclease-free water.

TEVC measurements were performed on cRNA- or water-injected oocytes (incubation time 3–5 days). The membrane voltage was clamped to -60 mV using a TEVC amplifier (Warner Instruments, Hamden, CT, USA), and transmembrane currents (I_M) were low-pass filtered at 1000 Hz (Frequency Devices 902, Haverhill, MA, USA) and recorded with a strip chart recorder (Kipp and Zonen, Delft, Netherlands). Oocytes

were placed in a perfusion chamber and perfused (gravity driven) with ORi without pyruvate and antibiotics (pH 7.4). In all experimental groups, measurements were performed on oocytes isolated from at least two different *Xenopus laevis* individuals.

For experiments examining the inhibition and recovery from inhibition of ACh-gated currents by LPC and G-PC, oocytes were injected with a 1:1 ratio of cRNA encoding human α 9 and α 10 or α 7 alone and incubated at 17°C for 2–3 days prior to use. Oocytes were placed in a 30 μ l chamber and continuously perfused by gravity with a solution containing (in mM) 96 NaCl, 2.5 KCl, 1.8 CaCl₂ and 1 MgCl₂ (pH 7.4). Stock solutions of LPC or G-PC were prepared in water and diluted with perfusion solution to a final concentration of 1 μ M or 100 μ M, respectively. The membranes were clamped at a holding potential of -70 mV and stimulated with 1 s pulses of 100 μ M ACh once every 60 s until a steady-state baseline response was observed. Then, the perfusion solution was switched to one containing LPC or G-PC and the ACh-evoked responses were monitored for changes in amplitude and the data for inhibition of the ACh-evoked responses were normalized to three averaged control pulses, in the absence of LPC or G-PC, and analyzed with an exponential decay equation. The data for inhibition and recovery of inhibition by LPC or G-PC were best fit with single exponential equations.

Statistical Analysis and Data Processing


Data were analyzed using SPSS software (Munich, Germany) by Wilcoxon signed-rank test or by the non-parametric Kruskal-Wallis test, followed by the Mann-Whitney rank-sum test. A p value below 0.05 was considered as statistically significant and marked with *. Data were visualized using program Inkscape version 0.92 (Free and Open Source Software licensed under the GPL).

RESULTS

Single Gene Deletions of *Chrna7*, *Chrna9* or *Chrna10* Fully Abolish ACh-, Nic- or PC-Mediated Inhibition of BzATP-Induced IL-1 β Release by Mouse PBMCs

ACh and Nic, known canonical agonists of nAChRs, as well as PC have been reported to inhibit ATP-mediated IL-1 β release by human monocytes by a mechanism involving nAChRs (Hecker et al., 2015; Richter et al., 2016). To further determine the involvement of nAChR subunits α 7, α 9 and/or α 10 in this mechanism, PBMCs from WT mice and mice deficient in single nAChR subunit genes were investigated. PBMCs isolated from WT mice or from *Chrna7*, *Chrna9* or *Chrna10* gene-deficient mice spontaneously released low amounts of IL-1 β into the cell culture medium (Figures 1A–C). In response to stimulation with BzATP, an agonist of ATP receptor P2X7, the concentration of IL-1 β was increased (Figures 1A–C). ACh, Nic and PC significantly inhibited the BzATP-induced release of IL-1 β from WT PBMCs (Figures 1A–C, $p \leq 0.05$, $n \geq 4$). In sharp contrast to cells isolated from WT animals, ACh, Nic and PC did not change BzATP-induced release of IL-1 β from PBMCs obtained from *Chrna7* gene-deficient mice (Figure 1A, $n \geq 5$). Similarly,

¹<https://www.addgene.org/62276/>



the ACh- and Nic-mediated inhibition of IL-1 β release was fully abolished in PBMCs from *Chrna9* (Figure 1B, $n \geq 5$) or *Chrna10* (Figure 1C, $n \geq 4$) gene-deficient mice. Previously, we demonstrated that the inhibitory effect of PC is also abolished in

PBMCs deficient in *Chrna9* and *Chrna10* (Richter et al., 2016). Cell death, estimated by measurement of LDH release, remained below 9% in all experiments (data not shown).

The nAChR Subunits $\alpha 7$, $\alpha 9$ and $\alpha 10$ Are Necessary for ACh-, Nic- or PC-Mediated Inhibition of BzATP-Induced IL-1 β Release by U937 Cells

To corroborate the results obtained in mouse PBMCs and to confirm that the inhibitory effects of ACh, Nic and PC depend on the activation of nAChRs composed of $\alpha 7$, $\alpha 9$ and/or $\alpha 10$ subunits in human monocytic cells as well, a panel of nAChR antagonists was used. As expected, ACh, Nic or PC completely inhibited the IL-1 β release from human monocytic U937 cells primed with LPS and stimulated with BzATP (Figures 2A–C, $p \leq 0.05$, $n = 4$). In the presence of the general nAChR antagonist Mec (Philip et al., 2012), the inhibitory effect of ACh and PC was lost (Figures 2A,C). Similarly, the nAChR antagonists α -Bun and Stry, that preferentially inhibit nAChRs containing $\alpha 7$ or $\alpha 9$ subunits (McIntosh et al., 2009; Kudryavtsev et al., 2015), effectively blocked the inhibitory effects of ACh and PC. As published before, Nic was also ineffective in the presence of Mec, α -Bun or Stry (Hecker et al., 2015). In line with these observations, ArIB [V11L, V16D], a selective antagonist of $\alpha 7$ nAChRs (Whiteaker et al., 2007; Innocent et al., 2008), abolished the inhibitory effects of ACh and Nic (Figures 2A,B), whereas the effect of PC was strongly diminished but not fully inhibited (Figure 2C). Suppression of the activity of the nAChR subunit $\alpha 9$ by the treatment with the selective α -conotoxin RgIA4 (Vincler et al., 2006; Romero et al., 2017) abolished the inhibitory effect mediated by ACh or Nic (Figures 2A,B). It has been shown before, that the inhibitory effect of PC was prevented in the presence of RgIA4 (Richter et al., 2016). Of note, the application of nAChR antagonists alone had no effect on the IL-1 β release (Hecker et al., 2015; Richter et al., 2016).

To further confirm the importance of nAChR subunits $\alpha 7$, $\alpha 9$ and/or $\alpha 10$ in the ACh-, and PC-mediated inhibition of BzATP-induced IL-1 β secretion, the expression of these subunits was down-regulated in U937 cells by siRNA. As expected, ACh and PC fully inhibited IL-1 β release from U937 cells transfected with scrambled siRNA (Figures 3A,B, $p \leq 0.05$, $n = 4$), whereas silencing the expression of the $\alpha 7$ subunit blunted the inhibitory potential of ACh and PC in U937 cells primed with LPS and stimulated with BzATP (Figures 3A,B). Similarly, a reduced expression of $\alpha 9$ or $\alpha 10$ subunits blocked the inhibitory mechanism mediated by ACh (Figure 3A). Of importance, gene-silencing did not impair BzATP-mediated IL-1 β release (Figures 3A,B).

To control for the efficiency and specificity of subunit knock-down, the mRNA expression of nAChR subunits $\alpha 9$ and $\alpha 10$ was quantified by real-time RT-PCR (Figures 4A–D). As already reported (Hecker et al., 2015), transfection of siRNAs targeting the $\alpha 9$ or the $\alpha 10$ nAChR subunit resulted in a significant down-regulation of mRNA of subunit $\alpha 9$ (Figure 4A, $p \leq 0.05$, $n = 4$) and $\alpha 10$ (Figure 4C, $p \leq 0.05$, $n = 4$) respectively, when compared with cells transfected with scrambled siRNA.




FIGURE 2 | Secretion of IL-1 β by U937 cells in the presence of nicotinic acetylcholine receptor (nAChR) antagonists and ACh, Nic or PC. U937 cells were primed with lipopolysaccharide (LPS; 1 μ g/ml) for 5 h and BzATP (100 μ M) was given for additional 30 min, in the presence or absence of ACh (10 μ M) (**A**), Nic (100 μ M) (**B**) or PC (100 μ M) (**C**). BzATP induced the release of IL-1 β , which was inhibited in the presence of ACh, Nic or PC (**A–C**). The inhibitory potential of ACh (**A**) was abolished by mecamylamine (Mec, 100 μ M), α -bungarotoxin (α -Bun, 1 μ M), strychnine (Stry, 10 μ M), ArlB [V11L, V16D] (500 nM) or RglA4 (200 nM). Addition of ArlB or RglA4 antagonized the effects of Nic (**B**). Similarly the effect of PC was blunted by Mec, α -Bun, Stry and ArlB (**C**). The IL-1 β concentration was quantified by ELISA. Data were analyzed by Kruskal-Wallis test followed by Mann-Whitney rank sum test and presented as individual data points, bars represent median, whiskers percentiles 25 and 75. * $p \leq 0.05$ significantly different from samples where BzATP was given alone.

In contrast, transfection with siRNA targeting nAChR subunits $\alpha 7$ or $\alpha 10$ did not change the $\alpha 9$ mRNA expression (**Figure 4B**). Similarly, transfection with siRNA specific for the $\alpha 7$ or $\alpha 9$ nAChR subunits did not influence the expression of subunit




FIGURE 3 | Down-regulation of nAChR subunits $\alpha 7$, $\alpha 9$ or $\alpha 10$ by siRNA abolished the inhibitory potential of ACh or PC. U937 cells were transfected with siRNA targeting *CHRNA7* ($\alpha 7$), *CHRNA9* ($\alpha 9$), *CHRNA10* ($\alpha 10$) or with scrambled siRNA. Forty-eight hours after transfection, cells were primed with lipopolysaccharide (LPS; 1 μ g/ml) for 5 h and BzATP (100 μ M) was given for additional 30 min in the presence or absence of ACh (10 μ M) (**A**) or PC (100 μ M) (**B**). BzATP induced the release of IL-1 β , which was inhibited in the presence of ACh or PC in cells transfected with scrambled siRNA. Down-regulation of the nAChR subunit $\alpha 7$ blunted the effect of ACh (**A**) or PC (**B**). Similarly, reduced expression of the $\alpha 9$ and $\alpha 10$ nAChR subunits diminished the inhibitory potential of ACh (**A**). The IL-1 β concentration was measured by ELISA. Data were analyzed by Kruskal-Wallis test followed by Mann-Whitney rank sum test and presented as individual data points, bars represent median, whiskers percentiles 25 and 75. * $p \leq 0.05$ significantly different from samples where BzATP was given alone.

$\alpha 10$ (**Figure 4D**). The mRNA expression of nAChR subunit $\alpha 7$ was detected at a very low level, which could not be reliably quantified. Due to the lack of specific antibodies (Moser et al., 2007; Rommel et al., 2015) it is problematic to investigate protein expression levels of nAChR subunits.

LPC and G-PC Dose-Dependently Inhibit BzATP-Induced IL-1 β Release by U937 Cells

To test whether other metabolites of phosphatidylcholines, besides PC, could also inhibit BzATP-induced IL-1 β release




FIGURE 4 | Reduction of mRNA expression of nAChR subunits $\alpha 9$ and $\alpha 10$ upon siRNA transfection. U937 cells were transfected with scrambled siRNA or with siRNA specific for *CHRNA7* ($\alpha 7$), *CHRNA9* ($\alpha 9$) or *CHRNA10* ($\alpha 10$). Forty-eight hours after transfection, the mRNA expression of $\alpha 9$ (A,B) or $\alpha 10$ (C,D) was analyzed by real-time RT-PCR. Transfection efficiently (A,C) and selectively (B,D) down-regulated the mRNA expression of the gene targeted with gene-specific siRNA. Data were analyzed by Mann-Whitney rank sum test and presented as individual data points, bars represent median, whiskers percentiles 25 and 75. * $p \leq 0.05$ significantly different from samples transfected with scrambled siRNA.

in U937 cells, the inhibitory effects of LPC and G-PC were examined. LPC ($IC_{50} \sim 1 \mu M$; **Figure 5A**, $p \leq 0.05$, $n = 5$) as well as G-PC ($IC_{50} \sim 1 \mu M$; **Figure 5B**, $p \leq 0.05$, $n \geq 4$) dose-dependently inhibited BzATP-induced IL-1 β release without affecting the viability of the cells, as measured by LDH release (data not shown). Application of LPC (100 μM) or G-PC (100 μM) alone did not induce IL-1 β release.


The nAChR Subunits $\alpha 9$ and $\alpha 10$ Are Mandatory for LPC- or G-PC-Mediated Inhibition of BzATP-Induced IL-1 β Release by U937 Cells

To analyze the involvement of nAChR subunits $\alpha 7$, $\alpha 9$ and/or $\alpha 10$ in the LPC- and G-PC-mediated inhibition of BzATP-induced IL-1 β secretion, a panel of nAChR antagonists was applied. The inhibitory effects of LPC and G-PC were fully prevented in the presence of Mec, α -Bun, Stry or RgIA4. In contrast, ArIB, the nAChR $\alpha 7$ specific antagonist, at best slightly attenuated the inhibitory effects mediated by LPC and G-PC (**Figures 6A,B**, $p \leq 0.05$, $n = 4$). To further confirm these results, the expression of nAChR subunits $\alpha 7$, $\alpha 9$ or $\alpha 10$ was selectively down-regulated

using siRNA (**Figures 7A,B**, $p \leq 0.05$, $n = 4$). Silencing of single nAChR subunits, or transfection of scrambled siRNA, did not influence the BzATP-induced IL-1 β release. Importantly, silencing the expression of nAChR subunits $\alpha 9$ or $\alpha 10$ effectively abolished the inhibitory potential of LPC and G-PC. In contrast, down-regulation of the $\alpha 7$ nAChR subunit only slightly impaired the inhibitory effects of LPC and G-PC (**Figures 7A,B**, $p \leq 0.05$, $n = 4$).


LPC and G-PC Do Not Induce Ion Channel Functions at Heterologously Expressed nAChRs

Next, we investigated the effect of LPC and G-PC at canonical ionotropic nAChRs. For this purpose, we used *Xenopus laevis* oocytes as a heterologous expression system for different combinations of human $\alpha 7$, $\alpha 9$ and $\alpha 10$ nAChR subunits and performed TEVC measurements to assess ion channel functions. We demonstrated previously, that PC does not evoke ion currents at heterologously expressed homomeric $\alpha 9$ and heteromeric $\alpha 9\alpha 10$ nAChR (Richter et al., 2016). Here we tested the effects of PC at different combinations of human $\alpha 7$, $\alpha 9$ and




$\alpha 10$ nAChR subunits. To control for the expression of nAChR in all experiments, the nAChR agonist ACh (100 μ M; Papke et al., 1996; Azam and McIntosh, 2012) was used as a positive control.

The first set of experiments was performed on oocytes that were injected with cRNA encoding for $\alpha 7$, $\alpha 9$ and $\alpha 10$ nAChR subunits (Figure 8). As expected, application of ACh (ACh1) evoked currents (ΔI_M) that were reversible and repeatable by subsequent ACh application (ACh2; Figure 8A). The ACh-induced effects (ACh1, ACh2) were not significantly different ($n = 17$, $p = 0.14$; Figures 8A,E). Next, LPC (100 μ M) was tested on oocytes expressing nAChR subunits $\alpha 7$, $\alpha 9$ and $\alpha 10$ (Figure 8B). While LPC (2 min) did not elicit ion currents, subsequent application of ACh resulted in significant current responses ($n = 8$, $p = 0.001$; Figures 8B,E). In the same way, G-PC (100 μ M; Figure 8C) and PC (1 mM; Figure 8D) were tested. Again, neither G-PC nor PC induced ion currents, while application of ACh induced a current response (G-PC: $n = 6$, $p = 0.001$, Figures 8C,E; PC: $n = 8$, $p = 0.001$; Figures 8D,E).




In water-injected control oocytes, ACh ($n = 4$), LPC ($n = 6$), G-PC ($n = 4$) and PC ($n = 6$) and had no impact on ΔI_M (Figures 8F–I).

In addition, we performed experiments on oocytes that were injected with different combinations of human nAChR subunits (Figure 9). In oocytes expressing homomeric $\alpha 7$ nAChRs, application of LPC, G-PC or PC had no impact on ΔI_M , while ACh induced significant current responses (Figures 9A,B; LPC: $n = 6$, $p = 0.001$; G-PC: $n = 6$, $p = 0.001$; PC: $n = 6$, $p = 0.001$). The same results were obtained in experiments on oocytes co-expressing nAChR subunits $\alpha 7$ and $\alpha 9$ (LPC: $n = 6$, $p = 0.001$; G-PC: $n = 7$, $p = 0.001$; PC: $n = 6$, $p = 0.001$; Figures 9C,D). It was previously



shown, that PC does not induce ion channel functions on heterologously expressed heteromeric $\alpha 9\alpha 10$ nAChR (Richter et al., 2016). Here, we investigated the impact of LPC and G-PC on heteromeric $\alpha 9\alpha 10$ nAChR (Figures 9E,F). Again, application of LPC and G-PC had no effect on ΔI_M , while ACh induced a significant current response (Figures 9E,F; LPC: $n = 6$, $p = 0.001$; G-PC: $n = 6$, $p = 0.001$).



In all experiments the ACh-induced effect was repeatable (ACh1, ACh2) and not significantly different compared to the first response ($\alpha 7$: $n = 9$, $p = 0.432$; $\alpha 7\alpha 9$: $n = 7$, $p = 0.645$; $\alpha 9\alpha 10$: $n = 6$, $p = 0.65$; Figures 9B,D,F). LPC, G-PC, PC and ACh did not induce changes in ion currents of water-injected control oocytes (data not shown).




FIGURE 9 | LPC, G-PC and PC do not induce ion channel activity at heterologously expressed nAChR subunits. TEVC measurements were performed on *Xenopus laevis* oocytes that expressed the human $\alpha 7$ nAChR subunit alone (**A,B**), as well as co-expressed human $\alpha 7$ and $\alpha 9$ (**C,D**) or human $\alpha 9$ and $\alpha 10$ (**E,F**) nAChR subunits. In all experiments, application of ACh (100 μ M; 1 s) induced repetitive stimulations of the transmembrane ion current (ΔI_M ; ACh1 and ACh2). In contrast, initial application of LPC (100 μ M; 2 min), G-PC (100 μ M; 2 min) as well as PC (100 μ M; 2 min) had no impact on I_M , whereas ACh induced a current response. All changes of ΔI_M induced by cholinergic stimulation are shown as individual data points, bars represent median, whiskers percentiles 25 and 75. Statistical analyses were performed using the Wilcoxon signed-rank test. * $p \leq 0.05$ significantly different from the ΔI_M values before.

Inhibition of ACh-Induced Currents by LPC and G-PC

Previously, we demonstrated that PC blunts classical ion channel responses at $\alpha 9\alpha 10$ nAChR heterologously expressed by *Xenopus laevis* oocytes (Richter et al., 2016). To test if LPC and G-PC have an impact on ACh-induced ion currents of nAChR, we performed TEVC measurements and monitored current responses induced by ACh (100 μ M, 1 s pulses) in presence or absence of LPC or G-PC (Figure 10). In experiments on oocytes expressing heteromeric $\alpha 9\alpha 10$ nAChR, LPC (1 μ M) decreased the ACh-evoked responses to $69 \pm 2\%$ ($n = 12$) of control values after a 12 min exposure to the compound (Figures 10A,B). Thereafter, LPC was washed out and the current responses recovered to $101 \pm 2\%$ ($n = 12$) after 12 min (Figures 10A,B). Similar experiments were performed on oocytes expressing homomeric $\alpha 7$ nAChR (Figure 10C). In presence of LPC, the ACh-evoked currents were reduced to $89 \pm 7\%$ ($n = 6$) of control values after a 12 min perfusion (Figure 10C). The responses recovered to $101 \pm 6\%$ ($n = 6$) of control values after a 12 min wash-out period (Figure 10C). In oocytes that expressed heteromeric $\alpha 9\alpha 10$ nAChR, G-PC (100 μ M) decreased the ACh-induced currents to $60 \pm 7\%$ ($n = 6$) of control values (Figure 10D). After a 12 min wash-out period of G-PC, the ACh-induced responses recovered to $90 \pm 2\%$ ($n = 6$) of control values (Figure 10D).

DISCUSSION

Anti-inflammatory effects of nAChR agonists in macrophages were repeatedly described and the term “anti-inflammatory cholinergic reflex” was coined more than a decade ago. It was suggested that ACh of vagal origin activates nAChR composed of subunit $\alpha 7$ and controls the synthesis of pro-IL-1 β and other pro-inflammatory cytokines on the transcriptional and post-transcriptional level (Borovikova et al., 2000; de Jonge et al., 2005; Rosas-Ballina et al., 2011; Olofsson et al., 2012). We recently discovered that Nic, Cho and ACh efficiently reduce the ATP-induced inflammasome activation and IL-1 β maturation in human monocytes. Ligand-binding to monocytic nAChRs containing subunits $\alpha 7$, $\alpha 9$ and/or $\alpha 10$ triggers a metabotropic response that efficiently inhibits ATP-signaling at P2X7 receptor (Hecker et al., 2015). Similar cholinergic effects on monocytes are also provoked by PC, PC-modified macromolecules and DPPC (Hecker et al., 2015; Richter et al., 2016; Backhaus et al., 2017). In the current study: (i) we demonstrate that LPC and G-PC, common metabolites of phosphatidylcholines, also activate monocytic nAChR; (ii) we determine the nAChR subunit requirements of different conventional and non-conventional nicotinic agonists; and (iii) we provide evidence that LPC, G-PC and PC do not evoke ion currents at heterologously expressed nAChR, but rather act as silent agonists.

We provide evidence that LPC and G-PC function as nAChR agonists that dose-dependently and efficiently inhibit the BzATP-mediated IL-1 β release by LPS-primed human monocytic U937 cells. Interestingly, the IC₅₀ of both

compounds is one order of magnitude lower than that of PC (Hecker et al., 2015) or DPPC (Backhaus et al., 2017). Importantly, effective concentrations of LPC, G-PC and PC can be detected in human plasma (Icol et al., 2002, 2005; Drobnik et al., 2003; Quehenberger et al., 2010). As LPC and G-PC are metabolites of phosphatidylcholines, different cell signaling pathways involving phospholipases as well as exposure to insect or snake venoms rich in phospholipases (Harris and Scott-Davey, 2013) may trigger this anti-inflammatory pathway. There are some conflicting data on immune-modulatory functions of LPC (Liu-Wu et al., 1998; Kabarowski et al., 2002; Stock et al., 2006; Carneiro et al., 2013), but none of these studies suggested a cholinergic mechanism. Anti-inflammatory effects of LPC, including a reduction of peritoneal IL-1 β levels, were reported for experimental sepsis in the mouse and these effects seemed to depend on the presence of 18 carbon molecules in the acyl chain (Yan et al., 2004), whereas we used LPC with a chain length of 16 in our experiments. Other studies suggested an induction of pro-IL-1 β in human monocytes during long-term incubation with LPC with a carboxyl chain length of 16 and 18 (Liu-Wu et al., 1998) and a LPC- and Ca²⁺-dependent induction of IL-1 β in LPS-primed microglial cells (Stock et al., 2006). These conflicting results might be explained by the different experimental protocols, the different cell types investigated and by the potential presence of pyrogenic contaminations that induce the expression of IL-1 β .

In order to define which nAChR subunits are required for the inhibitory function of ACh, Nic and PC on the BzATP-driven release of IL-1 β , we used gene-deficient mice, a panel of nAChR antagonists and performed gene-silencing experiments. We knew from previous pharmacological and gene-silencing studies that the effect of Nic is mediated via nAChR containing subunits $\alpha 7$, $\alpha 9$ and/or 10 (Hecker et al., 2015). It remained, however, unclear, if all three receptor subunits are mandatory or if they exert redundant functions. Regarding the signaling of PC, experiments on gene-deficient mice as well as pharmacological and gene-silencing studies revealed that nAChR subunits $\alpha 9$ and $\alpha 10$ are mandatory (Richter et al., 2016) but in this case the role of subunit $\alpha 7$ was still unclear and nothing was known about the nAChR subunit requirements for the ACh mediated effects. First, we investigated PBMCs from WT mice and mice deficient in the genes for nAChR subunits $\alpha 7$, $\alpha 9$ or $\alpha 10$. The results of these experiments suggested that all three receptor subunits are mandatory for the inhibition of IL-1 β release by ACh, Nic and PC. In the same line, Mec, a general nicotinic antagonist (Philip et al., 2012) as well as α -Bun and Stry, antagonists of nAChR containing subunits $\alpha 7$ and $\alpha 9$ (McIntosh et al., 2009; Kudryavtsev et al., 2015), efficiently reverted the inhibitory effects of ACh, Nic (Hecker et al., 2015) and PC in LPS-primed human U937 cells. Similarly, RgIA4, a selective antagonist of nAChR subunit $\alpha 9$ (Vincler et al., 2006; Romero et al., 2017) and ArIB, a selective antagonist of nAChR subunit $\alpha 7$ (Innocent et al., 2008), fully abolished the inhibitory effect mediated by ACh, Nic and PC (Richter et al., 2016). These data were further corroborated by siRNA experiments performed here and in previous studies (Hecker et al., 2015; Richter et al., 2016). We conclude from these data that all three nAChR subunits $\alpha 7$,




FIGURE 10 | LPC and G-PC desensitize ACh-gated currents mediated by nAChRs heterologously expressed in *Xenopus laevis* oocytes. TEVC experiments were performed on oocytes that expressed human homomeric $\alpha 7$ or heteromeric $\alpha 9\alpha 10$ nAChRs. **(A)** Representative traces of ACh-gated currents mediated by $\alpha 9\alpha 10$ nAChRs illustrating the inhibitory effect of LPC (1 μ M). The current traces represent 30 s recordings each and are shown concatenated omitting the 30 s gap between each individual trace. The oocytes were continuously perfused with control solution and stimulated with 1 s pulses of ACh (100 μ M) once per min until steady-state baseline responses were observed (indicated by the arrow). Subsequently, the control solution was switched to one containing LPC and the ACh-gated currents were monitored for changes in amplitude for 12 min. Thereafter, the perfusion solution was switched back to control and the ACh-gated currents were monitored for recovery. **(B)** Graphical representation of the experimental results shown in **(A)** ($n = 12$). **(C)** Similar experiments as those in **(A,B)** were performed on $\alpha 7$ nAChR expressing oocytes to assess the effects of LPC on ACh-gated currents mediated by this nAChR subtype ($n = 6$). **(D)** Analysis of inhibition and recovery from inhibition of ACh-gated currents by G-PC (100 μ M) on $\alpha 9\alpha 10$ nAChR expressing oocytes ($n = 6$). The error bars in all graphs denote the standard error of mean (SEM) from the indicated numbers of oocytes. The data in **(B–D)** were analyzed with single exponential equations and the X-axes represent the temporal duration of each experimental condition (absence or presence of LPC or G-PC).






$\alpha 9$ and $\alpha 10$ are needed for signaling of ACh, Nic and PC in monocytic cells (**Table 1**).

Next, we investigated the receptor requirements for the more complex compounds LPC and G-PC by using both nAChR antagonists and gene-silencing of nAChR subunits by siRNA in human U937 cells. In contrast to ACh, Nic and PC, we provide evidence that only nAChR subunits $\alpha 9$ and $\alpha 10$ are mandatory for the inhibitory mechanism mediated by LPC and G-PC. There might be a biologically insignificant contribution of subunit $\alpha 7$ because both gene silencing and Ar1B seemed to provoke a subtle impairment of the inhibitory effects of LPC and G-PC. This discrepancy might be explained by differences in the structure and the physicochemical properties among these compounds

(**Table 1**). Interestingly, the recently reported inhibitory effects of DPPC on ATP-driven inflammasome activation also strictly depended on the $\alpha 9$ subunit, but in contrast to LPC and G-PC, the $\alpha 10$ subunit is not mandatory but it can be functionally replaced by subunit $\alpha 7$ (Backhaus et al., 2017).

With all due caution, we propose that the $\alpha 9$ nAChR subunit is the critical subunit involved in the metabotropic regulation of inflammasome activation, irrespective of the structure of the nicotinic agonist. The $\alpha 9$ nAChR subunit interacts with $\alpha 7$ and/or $\alpha 10$ subunits depending on the complexity of the nAChR agonist. Small agonists such as ACh or PC need all three subunits $\alpha 7$, $\alpha 9$ and $\alpha 10$ for signaling, the agonists LPC and G-PC depend on $\alpha 9$ and $\alpha 10$, without an essential contribution

TABLE 1 | Concentration of agonists causing 50% inhibition (IC_{50}) of the adenosine triphosphate (ATP)-mediated release of interleukin-1 β (IL-1 β) in monocytes and the nAChR composition needed for the signaling.

Endogenous agonist	Approximate IC_{50}	Required nAChR subunits
ACh	10 μ M (Hecker et al., 2015)	$\alpha 7$, $\alpha 9$, $\alpha 10$
		
PC	10 μ M (Hecker et al., 2015)	$\alpha 7$, $\alpha 9$, $\alpha 10$
		
LPC	1 μ M	$\alpha 9$, $\alpha 10$
		
G-PC	1 μ M	$\alpha 9$, $\alpha 10$
		
DPPC	10 μ M (Backhaus et al., 2017)	$\alpha 9$ and $\alpha 7$ or $\alpha 10$
		

ACh, acetylcholine; DPPC, dipalmitoylphosphatidylcholine; G-PC, glycerophosphocholine; LPC, lysophosphatidylcholine; nAChR, nicotinic ACh receptor; R, fatty acid residue C16:0.

of subunit $\alpha 7$ (Table 1). The more bulky and hydrophobic phosphatidylcholines like DPPC, signal alternatively via subunit combination $\alpha 7\alpha 9$ or via $\alpha 9\alpha 10$ (Backhaus et al., 2017; Table 1). We have to admit that although this generalization is tempting, we only tested LPC (16:0) and DPPC up to now, but different variants of LPC and phosphatidylcholines have to be investigated before establishing a principle, and these are many and varied.

The following experiments were designed to clarify if LPC, G-PC and PC interact with canonical nAChRs that function as ligand-gated ion channels. We have previously shown that PC does not evoke ion currents in *Xenopus laevis* oocytes expressing human $\alpha 9$ or $\alpha 9\alpha 10$ nAChRs. Here, we investigated the short-term effect of LPC, G-PC and PC on ionotropic nAChR functions using *Xenopus laevis* oocytes as a heterologous expression system for human $\alpha 7$ nAChR as well as the expression of combinations of $\alpha 7$, $\alpha 9$ and $\alpha 10$ subunits. Due to the lack of specific antibodies to the nAChR investigated (Moser et al., 2007; Rommel et al., 2015), we were not able to test if all nAChR subunits were co-expressed in our experimental setup, but we were able to demonstrate that ACh induced characteristic ion currents in oocytes injected with cDNA for nAChR but not in control oocytes. In addition, we do not know which homomeric or heteromeric nAChR species form in co-injected oocytes. Of note, LPC, G-PC or PC did not induce ion currents in any of

the experimental settings, whereas the classical nAChR agonist ACh did.

Although PC does not induce ion currents at classical nAChR, it seems to function as a silent agonist that desensitizes $\alpha 9\alpha 10$ nAChR (Richter et al., 2016). Hence, we investigated the long-term effects of LPC and G-PC on heterologously expressed nAChR. While the brief application of 100 μ M LPC did not evoke responses in oocytes expressing nAChRs, application of LPC (100 μ M) induced a non-specific increase of the baseline holding current during continuous perfusion of the compound. Such effects have been described before and might be due to a detergent-like action of LPC (Ikeuchi et al., 1997). Therefore, we lowered the LPC concentration in the long-term experiments to 1 μ M, a concentration that still inhibited the ATP-induced IL-1 β release in U937 cells.

In the inhibition and recovery from inhibition measurements, we found that in human $\alpha 9\alpha 10$ nAChR-expressing oocytes, ACh-mediated current responses were blunted in the presence of LPC and G-PC. The slow kinetics of inhibition and recovery from inhibition by LPC and G-PC suggest that both compounds might also function as silent agonists of the heteromeric $\alpha 9\alpha 10$ nAChR. However, we cannot formally exclude that LPC and G-PC act as partial antagonists at these nAChR. Interestingly, while LPC decreased the ACh-evoked current responses to $69 \pm 2\%$ in heteromeric $\alpha 9\alpha 10$ nAChR expressing oocytes, the ACh-evoked currents in homomeric $\alpha 7$ nAChR expressing oocytes were decreased only to $89 \pm 7\%$. These findings are in accordance with our observation on the monocytic U937 cells, in which the immuno-modulatory function of LPC requires the expression of nAChR subunits $\alpha 9$ and $\alpha 10$ but not $\alpha 7$. In summary, LPC, G-PC and PC are potent agonists of non-classical nicotinic receptors of monocytes that provoke metabotropic functions. LPC, G-PC and PC do not induce ion currents, but they seem to act as silent agonists.

Receptor subunits $\alpha 7$, $\alpha 9$ and $\alpha 10$ belong to a unique class of evolutionary conserved nAChR, which in contrast to all other nAChR can form pentameric homomers (Franchini and Elgoyhen, 2006; Lipovsek et al., 2012, 2014). Homomeric $\alpha 7$ and $\alpha 9$ nAChR were also described (Elgoyhen et al., 1994), whereas the human $\alpha 10$ subunit alone does not form functional receptors (Elgoyhen et al., 2001). The $\alpha 9$ and $\alpha 10$ nAChR subunits can co-assemble to heteromers with pharmacological properties different from $\alpha 9$ homomers (Boffi et al., 2017). Heteromers of $\alpha 7$ and $\alpha 10$ have been evidenced in rat sympathetic neurons (Lips et al., 2006). Furthermore a complex interaction of nAChR subunits $\alpha 7$, $\alpha 9$ or $\alpha 10$ in regulating vestibular afferent gain was shown (Morley et al., 2017). In a mast cell line, interaction of nAChR subunits $\alpha 7$, $\alpha 9$ and $\alpha 10$ was described in the context of nicotinic inhibition of Fc ϵ RI-induced leukotriene and cytokine production (Mishra et al., 2010). However, exceedingly low nanomolar concentrations of Nic were effective in this mast cell line, which is not the case for the inhibition of ATP-induced release of IL-1 β by human U937 cells (Hecker et al., 2015). In all leukocyte subtypes tested so far, nAChR subunits do not form conventional ligand-gated ion channels,

but rather exert metabotropic function (Peng et al., 2004; Razani-Boroujerdi et al., 2007; Hecker et al., 2009, 2015; Mishra et al., 2010; Richter et al., 2016; Valbuena and Lerma, 2016). We published before, that Nic, Cho, PC and DPPC exert their immuno-modulatory functions via metabotropic inhibition of P2X7 receptor function (Hecker et al., 2015; Richter et al., 2016; Backhaus et al., 2017). However, the structure of non-canonical nAChR receptors exerting metabotropic functions is largely unknown; it is even unclear, if leukocytic nAChR form multimers.

There is increasing evidence, that signaling via nAChR subunit $\alpha 7$ induces Ca^{2+} signals and that signaling-related proteins including G proteins can interact with a cytoplasmic loop of this subunit (Kabbani et al., 2013; King and Kabbani, 2016; Valbuena and Lerma, 2016). Similar protein-protein interactions might also occur for subunit $\alpha 9$. In addition, direct interactions of nAChR containing subunit $\alpha 6$ with purinergic receptors P2X2 and P2X3 have been reported (Limapichat et al., 2014; Wieskopf et al., 2015). This interesting and emerging field of metabotropic nicotinic signaling certainly deserves more scientific attention.

Excessive release of IL-1 β into the circulation in response to trauma-associated extracellular ATP is useless and dangerous, because the cytokine is swept away from the site of injury and can cause severe life-threatening SIRS and multi-organ damage (Stoecklein et al., 2012; Lord et al., 2014). As monocytes are the major source of IL-1 β within the circulation (Arango Duque and Descoteaux, 2014), the inhibitory mechanisms described in this study might be of high clinical relevance. ATP-driven inflammasome activation is typical for the undesired sterile inflammation that causes a high morbidity and mortality worldwide, whereas several ATP-independent pathways lead to the desired inflammasome activation needed for defense against infection (Cauwels et al., 2014). Of note, nAChR subunit $\alpha 9$ plays a central role in the control of ATP-driven inflammasome activation in monocytes, whereas subunit $\alpha 7$ controls the expression of pro-inflammatory cytokines more generally (Borovikova et al., 2000). The specific $\alpha 9$ -dependend pathway seems to be an interesting target that might enable the prevention of SIRS caused by multiple traumata and major surgery without inhibiting host defense against infections. As we demonstrated that the cholinergic inhibition of ATP-driven IL-1 β release is also active in mice, this concept can and should be evaluated experimentally *in vivo*.

This study has several limitations and more research is needed to further corroborate our concept. One major technical issue is the lack of specific antibodies to the nAChR subunits under investigation that prevents an unequivocal proof of the specificity and efficiency of gene-silencing as well as the proof of the heterologous co-expression of nAChR subunits. At present, we cannot conclude that all LPC species act in the same way like LPC (16:0) and the same holds true for phosphatidylcholines, where only DPPC was investigated previously (Backhaus et al., 2017). It also remains to be tested, if LPC and G-PC stimulate ion currents in monocytic cells, although this is improbable as PC and DPPC do not (Richter et al., 2016; Backhaus et al., 2017). Additionally, more

research is needed to identify the signaling pathways involved in the inhibitory mechanism mediated by LPC, G-PC and PC. Finally, we can only speculate on the relevance of our findings *in vivo*.

In conclusion, we demonstrated that canonical nAChR agonists as well as PC inhibit ATP-induced release of IL-1 β by human and mouse mononuclear cells via nAChR subunits $\alpha 7$, $\alpha 9$ and $\alpha 10$. In contrast, LPC (16:0) and G-PC, more complex metabolites of phosphatidylcholines, elicit the inhibitory mechanism in human monocytes via interaction of the two nAChR subunits $\alpha 9$ and $\alpha 10$. Independent of the complexity of the compound, neither LPC, G-PC nor PC induce ionotropic functions at heterologously expressed nAChR, suggesting that PC-bearing molecules in general regulate immuno-modulatory functions without inducing canonical ion channel function at nAChR. This selective activity opens the opportunity for the development of novel anti-inflammatory therapies without inducing ionotropic functions of nAChRs.

AUTHOR CONTRIBUTIONS

AZ and KR participated in research design, performance of experiments, interpretation of the data and writing of the manuscript. AA, SW, KS, BF and AJH: performance of experiments, interpretation of the data and editing of the manuscript. GK-C, MA and WP participated in research design, interpretation of the data and editing of the manuscript. JMM and VG participated in research design, interpretation of the data and writing of the manuscript.

FUNDING

This study was supported by the LOEWE Program of the State of Hessen (Non-neuronal cholinergic systems) to VG, research grant from the German Research Foundation to VG (GR 1094/7-1). GK-C received funding by SFB TR84. JMM received funding from NIH Grants P01-GM48677 and R01-GM103801.

ACKNOWLEDGMENTS

We thank Sabine Stumpf and Kathrin Petri (Department of General and Thoracic Surgery, Laboratory of Experimental Surgery, Justus-Liebig-University Giessen) for excellent technical support. Lukas Wichmann (Institute for Animal Physiology, Justus-Liebig-University of Giessen) and Ivan Manzini (Institute for Animal Physiology and Molecular Biomedicine, Justus-Liebig-University of Giessen) for their help and support. We are grateful to Douglas E. Vetter (University of Mississippi Medical Center, Department of Neurobiology and Anatomical Sciences, Jackson, MS, USA) for providing the *Chrna9* and *Chrna10* gene-deficient mice, as well as Günther Schmalzing and Ralf Hausmann (Department of Molecular Pharmacology, RWTH Aachen University, Aachen, Germany) for providing cRNA encoding for human $\alpha 7$ nAChR.

REFERENCES

- Arango Duque, G., and Descoteaux, A. (2014). Macrophage cytokines: involvement in immunity and infectious diseases. *Front. Immunol.* 5:491. doi: 10.3389/fimmu.2014.00491
- Azam, L., and McIntosh, J. M. (2012). Molecular basis for the differential sensitivity of rat and human $\alpha 9\alpha 10$ nAChRs to α -conotoxin RgIA. *J. Neurochem.* 122, 1137–1144. doi: 10.1111/j.1471-4159.2012.07867.x
- Backhaus, S., Zakrzewicz, A., Richter, K., Damm, J., Wilker, S., Fuchs-Moll, G., et al. (2017). Surfactant inhibits ATP-induced release of interleukin-1 β ; via nicotinic acetylcholine receptors. *J. Lipid Res.* 58, 1055–1066. doi: 10.1194/jlr.M071506
- Boffi, J. C., Marcovich, I., Gill-Thind, J. K., Corradi, J., Collins, T., Lipovsek, M. M., et al. (2017). Differential contribution of subunit interfaces to $\alpha 9\alpha 10$ nicotinic acetylcholine receptor function. *Mol. Pharmacol.* 91, 250–262. doi: 10.1124/mol.116.107482
- Borovikova, L. V., Ivanova, S., Zhang, M., Yang, H., Botchkina, G. I., Watkins, L. R., et al. (2000). Vagus nerve stimulation attenuates the systemic inflammatory response to endotoxin. *Nature* 405, 458–462. doi: 10.1038/35013070
- Carneiro, A. B., Iaciura, B. M., Nohara, L. L., Lopes, C. D., Veas, E. M., Mariano, V. S., et al. (2013). Lysophosphatidylcholine triggers TLR2- and TLR4-mediated signaling pathways but counteracts LPS-induced NO synthesis in peritoneal macrophages by inhibiting NF- κ B translocation and MAPK/ERK phosphorylation. *PLoS One* 8:e76233. doi: 10.1371/journal.pone.0076233
- Cauwels, A., Rogge, E., Vandendriessche, B., Shiva, S., and Brouckaert, P. (2014). Extracellular ATP drives systemic inflammation, tissue damage and mortality. *Cell Death Dis.* 5:e1102. doi: 10.1038/cddis.2014.70
- de Jonge, W. J., van der Zanden, E. P., The, F. O., Bijlsma, M. F., van Westerloo, D. J., Bennink, R. J., et al. (2005). Stimulation of the vagus nerve attenuates macrophage activation by activating the Jak2-STAT3 signaling pathway. *Nat. Immunol.* 6, 844–851. doi: 10.1038/ni1229
- Dinarello, C. A., Simon, A., and van der Meer, J. W. (2012). Treating inflammation by blocking interleukin-1 in a broad spectrum of diseases. *Nat. Rev. Drug Discov.* 11, 633–652. doi: 10.1038/nrd3800
- Drobnik, W., Liebisch, G., Audebert, F. X., Frohlich, D., Gluck, T., Vogel, P., et al. (2003). Plasma ceramide and lysophosphatidylcholine inversely correlate with mortality in sepsis patients. *J. Lipid Res.* 44, 754–761. doi: 10.1194/jlr.M200401-jlr200
- Elgoyhen, A. B., Johnson, D. S., Boulter, J., Vetter, D. E., and Heinemann, S. (1994). $\alpha 9$: an acetylcholine receptor with novel pharmacological properties expressed in rat cochlear hair cells. *Cell* 79, 705–715. doi: 10.1016/0092-8674(94)90555-x
- Elgoyhen, A. B., Vetter, D. E., Katz, E., Rothlin, C. V., Heinemann, S. F., and Boulter, J. (2001). $\alpha 10$: a determinant of nicotinic cholinergic receptor function in mammalian vestibular and cochlear mechanosensory hair cells. *Proc. Natl. Acad. Sci. U S A* 98, 3501–3506. doi: 10.1073/pnas.051622798
- Ferrari, D., Pizzirani, C., Adinolfi, E., Lemoli, R. M., Curti, A., Idzko, M., et al. (2006). The P2X7 receptor: a key player in IL-1 processing and release. *J. Immunol.* 176, 3877–3883. doi: 10.4049/jimmunol.176.7.3877
- Franchini, L. F., and Elgoyhen, A. B. (2006). Adaptive evolution in mammalian proteins involved in cochlear outer hair cell electromotility. *Mol. Phylogenet. Evol.* 41, 622–635. doi: 10.1016/j.ympev.2006.05.042
- Froehner, S. C., Luetje, C. W., Scotland, P. B., and Patrick, J. (1990). The postsynaptic 43K protein clusters muscle nicotinic acetylcholine receptors in *Xenopus* oocytes. *Neuron* 5, 403–410. doi: 10.1016/0896-6273(90)90079-u
- Gloor, S., Pongs, O., and Schmalzing, G. (1995). A vector for the synthesis of cRNAs encoding Myc epitope-tagged proteins in *Xenopus laevis* oocytes. *Gene* 160, 213–217. doi: 10.1016/0378-1119(95)00226-v
- Gross, O., Thomas, C. J., Guarda, G., and Tschopp, J. (2011). The inflammasome: an integrated view. *Immunol. Rev.* 243, 136–151. doi: 10.1111/j.1600-065x.2011.01046.x
- Harris, J. B., and Scott-Davey, T. (2013). Secreted phospholipases A2 of snake venoms: effects on the peripheral neuromuscular system with comments on the role of phospholipases A2 in disorders of the CNS and their uses in industry. *Toxins (Basel)* 5, 2533–2571. doi: 10.3390/toxins5122533
- Hecker, A., Küllmar, M., Wilker, S., Richter, K., Zakrzewicz, A., Atanasova, S., et al. (2015). Phosphocholine-modified macromolecules and canonical nicotinic agonists inhibit ATP-induced IL-1 β release. *J. Immunol.* 195, 2325–2334. doi: 10.4049/jimmunol.1400974
- Hecker, A., Mikulski, Z., Lips, K. S., Pfeil, U., Zakrzewicz, A., Wilker, S., et al. (2009). Pivotal advance: up-regulation of acetylcholine synthesis and paracrine cholinergic signaling in intravascular transplant leukocytes during rejection of rat renal allografts. *J. Leukoc. Biol.* 86, 13–22. doi: 10.1189/jlb.1107722
- Ikeuchi, Y., Nishizaki, T., Matsuoka, T., and Sumikawa, K. (1997). Long-lasting enhancement of ACh receptor currents by lysophospholipids. *Brain Res. Mol. Brain Res.* 45, 317–320. doi: 10.1016/s0169-328x(96)00292-6
- Ilcol, Y. O., Dilek, K., Yurtkuran, M., and Ulus, I. H. (2002). Changes of plasma free choline and choline-containing compounds' concentrations and choline loss during hemodialysis in ESRD patients. *Clin. Biochem.* 35, 233–239. doi: 10.1016/s0009-9120(02)00298-9
- Ilcol, Y. O., Ozbek, R., Hamurtekin, E., and Ulus, I. H. (2005). Choline status in newborns, infants, children, breast-feeding women, breast-fed infants and human breast milk. *J. Nutr. Biochem.* 16, 489–499. doi: 10.1016/j.jnutbio.2005.01.011
- Innocent, N., Livingstone, P. D., Hone, A., Kimura, A., Young, T., Whiteaker, P., et al. (2008). α -conotoxin Arenatus IB[V11L,V16D] [corrected] is a potent and selective antagonist at rat and human native $\alpha 7$ nicotinic acetylcholine receptors. *J. Pharmacol. Exp. Ther.* 327, 529–537. doi: 10.1124/jpet.108.142943
- Kabarowski, J. H., Xu, Y., and Witte, O. N. (2002). Lysophosphatidylcholine as a ligand for immunoregulation. *Biochem. Pharmacol.* 64, 161–167. doi: 10.1016/s0006-2952(02)01179-6
- Kabbani, N., Nordman, J. C., Corgiat, B. A., Veltri, D. P., Shehu, A., Seymour, V. A., et al. (2013). Are nicotinic acetylcholine receptors coupled to G proteins? *Bioessays* 35, 1025–1034. doi: 10.1002/bies.201300082
- Kawashima, K., Fujii, T., Moriwaki, Y., Misawa, H., and Horiguchi, K. (2012). Reconciling neuronally and nonneuronally derived acetylcholine in the regulation of immune function. *Ann. N Y Acad. Sci.* 1261, 7–17. doi: 10.1111/j.1749-6632.2012.06516.x
- King, J. R., and Kabbani, N. (2016). Alpha7 nicotinic receptor coupling to heterotrimeric G proteins modulates RhoA activation, cytoskeletal motility, and structural growth. *J. Neurochem.* 138, 532–545. doi: 10.1111/jnc.13660
- Kudryavtsev, D., Shelukhina, I., Vulpius, C., Makarieva, T., Stonik, V., Zhmak, M., et al. (2015). Natural compounds interacting with nicotinic acetylcholine receptors: from low-molecular weight ones to peptides and proteins. *Toxins* 7, 1683–1701. doi: 10.3390/toxins7051683
- Limapichat, W., Dougherty, D. A., and Lester, H. A. (2014). Subtype-specific mechanisms for functional interaction between $\alpha 6\beta 4^*$ nicotinic acetylcholine receptors and P2X receptors. *Mol. Pharmacol.* 86, 263–274. doi: 10.1124/mol.114.093179
- Lipovsek, M., Fierro, A., Pérez, E. G., Boffi, J. C., Millar, N. S., Fuchs, P. A., et al. (2014). Tracking the molecular evolution of calcium permeability in a nicotinic acetylcholine receptor. *Mol. Biol. Evol.* 31, 3250–3265. doi: 10.1093/molbev/msu258
- Lipovsek, M., Im, G. J., Franchini, L. F., Pisciotto, F., Katz, E., Fuchs, P. A., et al. (2012). Phylogenetic differences in calcium permeability of the auditory hair cell cholinergic nicotinic receptor. *Proc. Natl. Acad. Sci. U S A* 109, 4308–4313. doi: 10.1073/pnas.1115488109
- Lips, K. S., König, P., Schätzle, K., Pfeil, U., Krasteva, G., Spies, M., et al. (2006). Coexpression and spatial association of nicotinic acetylcholine receptor subunits $\alpha 7$ and $\alpha 10$ in rat sympathetic neurons. *J. Mol. Neurosci.* 30, 15–16. doi: 10.1385/jmn.30:1.15
- Liu-Wu, Y., Hurt-Camejo, E., and Wiklund, O. (1998). Lysophosphatidylcholine induces the production of IL-1 β by human monocytes. *Atherosclerosis* 137, 351–357. doi: 10.1016/s0021-9150(97)00295-5
- Lord, J. M., Midwinter, M. J., Chen, Y. F., Belli, A., Brohi, K., Kovacs, E. J., et al. (2014). The systemic immune response to trauma: an overview of pathophysiology and treatment. *Lancet* 384, 1455–1465. doi: 10.1016/s0140-6736(14)60687-5
- McIntosh, J. M., Absalom, N., Chebib, M., Elgoyhen, A. B., and Vincler, M. (2009). $\alpha 9$ nicotinic acetylcholine receptors and the treatment of pain. *Biochem. Pharmacol.* 78, 693–702. doi: 10.1016/j.bcp.2009.05.020
- Mikulski, Z., Hartmann, P., Jositsch, G., Zaslon, Z., Lips, K. S., Pfeil, U., et al. (2010). Nicotinic receptors on rat alveolar macrophages dampen ATP-induced increase in cytosolic calcium concentration. *Respir. Res.* 11:133. doi: 10.1186/1465-9921-11-133

- Mishra, N. C., Rir-sima-ah, J., Boyd, R. T., Singh, S. P., Gundavarapu, S., Langley, R. J., et al. (2010). Nicotine inhibits Fc epsilon RI-induced cysteinyl leukotrienes and cytokine production without affecting mast cell degranulation through $\alpha 7/\alpha 10$ -nicotinic receptors. *J. Immunol.* 185, 588–596. doi: 10.4049/jimmunol.0902227
- Moosig, F., Damm, F., Knorr-Spahr, A., Ritgen, M., Zeuner, R. A., Kneba, M., et al. (2006). Reduced expression of C1q-mRNA in monocytes from patients with systemic lupus erythematosus. *Clin. Exp. Immunol.* 146, 409–416. doi: 10.1111/j.1365-2249.2006.03225.x
- Morley, B. J., Lysakowski, A., Vijayakumar, S., Menapace, D., and Jones, T. A. (2017). Nicotinic acetylcholine receptors regulate vestibular afferent gain and activation timing. *J. Comp. Neurol.* 525, 1216–1233. doi: 10.1002/cne.24131
- Moser, N., Mechawar, N., Jones, I., Gochberg-Sarver, A., Orr-Urtreger, A., Plomann, M., et al. (2007). Evaluating the suitability of nicotinic acetylcholine receptor antibodies for standard immunodetection procedures. *J. Neurochem.* 102, 479–492. doi: 10.1111/j.1471-4159.2007.04498.x
- Olofsson, P. S., Rosas-Ballina, M., Levine, Y. A., and Tracey, K. J. (2012). Rethinking inflammation: neural circuits in the regulation of immunity. *Immunol. Rev.* 248, 188–204. doi: 10.1111/j.1600-065x.2012.01138.x
- Orr-Urtreger, A., Göldner, F. M., Saeki, M., Lorenzo, I., Goldberg, L., De Biasi, M., et al. (1997). Mice deficient in the $\alpha 7$ neuronal nicotinic acetylcholine receptor lack α -bungarotoxin binding sites and hippocampal fast nicotinic currents. *J. Neurosci.* 17, 9165–9171.
- Ozaki, E., Campbell, M., and Doyle, S. L. (2015). Targeting the NLRP3 inflammasome in chronic inflammatory diseases: current perspectives. *J. Inflamm. Res.* 8, 15–27. doi: 10.2147/jir.s1250
- Papke, R. L., Bencherif, M., and Lippiello, P. (1996). An evaluation of neuronal nicotinic acetylcholine receptor activation by quaternary nitrogen compounds indicates that choline is selective for the $\alpha 7$ subtype. *Neurosci. Lett.* 213, 201–204. doi: 10.1016/s0304-3940(96)12889-5
- Peng, H., Ferris, R. L., Matthews, T., Hiel, H., Lopez-Albaitero, A., and Lustig, L. R. (2004). Characterization of the human nicotinic acetylcholine receptor subunit α (α) 9 (CHRNA9) and α (α) 10 (CHRNA10) in lymphocytes. *Life Sci.* 76, 263–280. doi: 10.1016/j.lfs.2004.05.031
- Philip, N. S., Carpenter, L. L., Tyrka, A. R., and Price, L. H. (2012). The nicotinic acetylcholine receptor as a target for antidepressant drug development. *ScientificWorldJournal* 2012:104105. doi: 10.1100/2012/104105
- Quehenberger, O., Armando, A. M., Brown, A. H., Milne, S. B., Myers, D. S., Merrill, A. H., et al. (2010). Lipidomics reveals a remarkable diversity of lipids in human plasma. *J. Lipid Res.* 51, 3299–3305. doi: 10.1194/jlr.M009449
- Ramarao, M. K., and Cohen, J. B. (1998). Mechanism of nicotinic acetylcholine receptor cluster formation by rapsyn. *Proc. Natl. Acad. Sci. U S A* 95, 4007–4012. doi: 10.1073/pnas.95.7.4007
- Rathinam, V. A., Vanaja, S. K., and Fitzgerald, K. A. (2012). Regulation of inflammasome signaling. *Nat. Immunol.* 13, 333–342. doi: 10.1038/ni.2237
- Razani-Boroujerdi, S., Boyd, R. T., Dávila-García, M. I., Nandi, J. S., Mishra, N. C., Singh, S. P., et al. (2007). T cells express $\alpha 7$ -nicotinic acetylcholine receptor subunits that require a functional TCR and leukocyte-specific protein tyrosine kinase for nicotine-induced Ca^{2+} response. *J. Immunol.* 179, 2889–2898. doi: 10.4049/jimmunol.179.5.2889
- Richter, K., Mathes, V., Fronius, M., Althaus, M., Hecker, A., Krasteva-Christ, G., et al. (2016). Phosphocholine—an agonist of metabotropic but not of ionotropic functions of $\alpha 9$ -containing nicotinic acetylcholine receptors. *Sci. Rep.* 6:28660. doi: 10.1038/srep28660
- Romero, H. K., Christensen, S. B., Di Cesare Mannelli, L., Gajewiak, J., Ramachandra, R., Elmslie, K. S., et al. (2017). Inhibition of $\alpha 9\alpha 10$ nicotinic acetylcholine receptors prevents chemotherapy-induced neuropathic pain. *Proc. Natl. Acad. Sci. U S A* 114, E1825–E1832. doi: 10.1073/pnas.1621433114
- Rommel, F. R., Raghavan, B., Paddenberg, R., Kummer, W., Tumala, S., Lochnit, G., et al. (2015). Suitability of nicotinic acetylcholine receptor $\alpha 7$ and muscarinic acetylcholine receptor 3 antibodies for immune detection: evaluation in murine skin. *J. Histochem. Cytochem.* 63, 329–339. doi: 10.1369/0022155415575028
- Rosas-Ballina, M., Olofsson, P. S., Ochani, M., Valdés-Ferrer, S. I., Levine, Y. A., Reardon, C., et al. (2011). Acetylcholine-synthesizing T cells relay neural signals in a vagus nerve circuit. *Science* 334, 98–101. doi: 10.1126/science.1209985
- Stock, C., Schilling, T., Schwab, A., and Eder, C. (2006). Lysophosphatidylcholine stimulates IL-1 β release from microglia via a P2X $_7$ receptor-independent mechanism. *J. Immunol.* 177, 8560–8568. doi: 10.4049/jimmunol.177.12.8560
- Stoecklein, V. M., Osuka, A., and Lederer, J. A. (2012). Trauma equals danger—damage control by the immune system. *J. Leukoc. Biol.* 92, 539–551. doi: 10.1189/jlb.0212072
- Stolz, M., Klapperstück, M., Kendzierski, T., Detro-Dassen, S., Panning, A., Schmalzing, G., et al. (2015). Homodimeric anoctamin-1, but not homodimeric anoctamin-6, is activated by calcium increases mediated by the P2Y $_1$ and P2X $_7$ receptors. *Pflugers Arch.* 467, 2121–2140. doi: 10.1007/s00424-015-1687-3
- Trautmann, A. (2009). Extracellular ATP in the immune system: more than just a “danger signal”. *Sci. Signal.* 2:pe6. doi: 10.1126/scisignal.256pe6
- Valbuena, S., and Lerma, J. (2016). Non-canonical signaling, the hidden life of ligand-gated ion channels. *Neuron* 92, 316–329. doi: 10.1016/j.neuron.2016.10.016
- Vetter, D. E., Katz, E., Maison, S. F., Taranda, J., Turcan, S., Ballesterio, J., et al. (2007). The $\alpha 10$ nicotinic acetylcholine receptor subunit is required for normal synaptic function and integrity of the olivocochlear system. *Proc. Natl. Acad. Sci. U S A* 104, 20594–20599. doi: 10.1073/pnas.0708545105
- Vetter, D. E., Liberman, M. C., Mann, J., Barhanin, J., Boulter, J., Brown, M. C., et al. (1999). Role of $\alpha 9$ nicotinic ACh receptor subunits in the development and function of cochlear efferent innervation. *Neuron* 23, 93–103. doi: 10.1016/s0896-6273(00)80756-4
- Vinler, M., Wittenauer, S., Parker, R., Ellison, M., Olivera, B. M., and McIntosh, J. M. (2006). Molecular mechanism for analgesia involving specific antagonism of $\alpha 9\alpha 10$ nicotinic acetylcholine receptors. *Proc. Natl. Acad. Sci. U S A* 103, 17880–17884. doi: 10.1073/pnas.0608715103
- Vladimer, G. I., Marty-Roix, R., Ghosh, S., Weng, D., and Lien, E. (2013). Inflammasomes and host defenses against bacterial infections. *Curr. Opin. Microbiol.* 16, 23–31. doi: 10.1016/j.mib.2012.11.008
- Wang, Y., Xiao, C., Indersmitten, T., Freedman, R., Leonard, S., and Lester, H. A. (2014). The duplicated $\alpha 7$ subunits assemble and form functional nicotinic receptors with the full-length $\alpha 7$. *J. Biol. Chem.* 289, 26451–26463. doi: 10.1074/jbc.M114.582858
- Whiteaker, P., Christensen, S., Yoshikami, D., Dowell, C., Watkins, M., Gulyas, J., et al. (2007). Discovery, synthesis, and structure activity of a highly selective $\alpha 7$ nicotinic acetylcholine receptor antagonist. *Biochemistry* 46, 6628–6638. doi: 10.1021/bi7004202
- Wieskopf, J. S., Mathur, J., Limapichat, W., Post, M. R., Al-Qazzaz, M., Sorge, R. E., et al. (2015). The nicotinic $\alpha 6$ subunit gene determines variability in chronic pain sensitivity via cross-inhibition of P2X $_2/3$ receptors. *Sci. Transl. Med.* 7:287ra272. doi: 10.1126/scitranslmed.3009986
- Wilczynska, J., Pfeil, U., Zakrzewicz, A., Dietrich, H., Körner, C., Hecker, A., et al. (2011). Acetylcholine and chronic vasculopathy in rat renal allografts. *Transplantation* 91, 263–270. doi: 10.1097/TP.0b013e3182038654
- Yan, J. J., Jung, J. S., Lee, J. E., Lee, J., Huh, S. O., Kim, H. S., et al. (2004). Therapeutic effects of lysophosphatidylcholine in experimental sepsis. *Nat. Med.* 10, 161–167. doi: 10.1038/nm989

Conflict of Interest Statement: Certain conotoxins, including RgIA4 have been patented by the University of Utah; JMM is an inventor on these patents.

The other authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2017 Zakrzewicz, Richter, Agné, Wilker, Siebers, Fink, Krasteva-Christ, Althaus, Padberg, Hone, McIntosh and Grau. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.